



UNIVERSITI PUTRA MALAYSIA

**CHROMOSOMAL REARRANGEMENT AND LOSS OF
HETEROZYGOSITY IN CERVICAL CANCER AMONGST PATIENTS
IN HOSPITAL KUALA LUMPUR**

SITI NORLASIAH BINTI ISMAIL.

IB 2004 2

**CHROMOSOMAL REARRANGEMENT AND LOSS OF
HETEROZYGOSITY STUDIES IN CERVICAL CANCER AMONGST
PATIENTS IN HOSPITAL KUALA LUMPUR**

By

SITI NORLASIAH BINTI ISMAIL

**Thesis Submitted to the School of Graduate Studies, Universiti
Putra Malaysia in Fulfilment of the Requirement for the Degree of
Doctor of Philosophy**

January 2004



For my beloved husband, Mat, my children, Nina, Niza, Amirah and Muneer, my mother, Siti Rokiah, my late father, Ismail, my brothers and sisters.

.....Allah will raise up, to (suitable) ranks (and degrees), those of you who believe and who have been granted knowledge.....

(Surah 58 Al - Mujaadalah : 11)

Abstract of thesis submitted to the senate of Universiti Putra Malaysia in fulfilment of the requirement for the degree of Doctor of Philosophy

**CHROMOSOMAL REARRANGEMENT AND LOSS OF
HETEROZYGOSITY STUDIES IN CERVICAL CANCER AMONGST
PATIENTS IN HOSPITAL KUALA LUMPUR**

By

SITI NORLASIAH BINTI ISMAIL

January 2004

Chairman: Professor Abdul Manaf Ali, Ph.D

Institute: Bioscience

Cervical carcinoma is the second most common malignancy among women worldwide. The highest incidence rates are observed in developing countries. The susceptibility to cervical carcinoma in high incidence populations may result from several factors including human papillomavirus (HPV) exposure and both inherited and acquired genes. HPV infection does not always led to cervical cancer. In cervical carcinoma the other common genetic characteristic of cancer is the presence of several recurrent genetic alterations, not related to HPV. The type of recurrent genetic damage might take different forms such as gene amplification, chromosomal translocation, deletions, loss of heterozygosity, point mutation, transcriptional silencing, and in some tumors viral DNA integration.

To determine the possible genetic alterations amongst the Malaysian women with cervical cancer, this study was conducted on 50 cervical



tumor biopsies received from the Department of Obstetrics and Gynecology of the Hospital Kuala Lumpur. The objectives of the study were; i) the establishment of short-term primary cell culture of human cervical epithelial cells derived from cervical tumors for the determination of the prevalence and the type of chromosomal aberrations, ii) characterization of the allelic losses of the chromosomes 3p, 5p, 11 and 17p (p53), subsequently identification of a possible site of candidate tumor suppressor gene(s) and iii) to determine the HPV status of the cervical cancers. Primary cell culture and cytogenetic techniques were performed on the cervical tumor biopsies. G-banding was employed for the identification of the chromosomes. To define the localization of the tumor suppressor genes, loss of heterozygosity study was performed on 37 cervical cancer cells. Twenty-four microsatellite polymorphic markers for the chromosomes 3p, 5p, 11 and p53 were chosen, the normal and tumor DNAs from each patient were analyzed for the allelic loss using PCR-based microsatellite analysis. The status for HPV 16 E6 and HPV 18 E6 was detected by PCR method.

Twenty-five cervical cancer biopsies were successfully karyotyped and near-diploid was the modal number, with a majority of them being hypodiploid (35-45). About 50% of the metaphases obtained in the 25 tumors were hypodiploids, 12.1% were hyperdiploids, and 36.7% were diploids. Numerical abnormalities were predominantly observed in the patients, with monosomies of chromosomes 17, 22, X, 11, 18, 19, 13, and 6. Fluorescence *in situ* hybridization using centromeric

probes 11, 17 and 18 confirmed the presence of monosomies 11, 17, 18 in a low percentage, 12.0%, 16.2% and 26.4% respectively. Several clones of cells were observed, with possibility of 45,XX,-22; 45,XX,-11; 45,XX,-19 and 45,XX,-18. Twenty-six of the 36 informative individuals exhibited LOH at one locus or more. The highest incidence was observed in chromosome 3p with the frequency of 48.6%, while a low frequency of 2.7% was detected in the short-arm of chromosome 17 at position 17p13.1, whereby lies the p53 tumor suppressor. LOH was confined to four prominent regions, 11q23.3, 3p14.2-3p14.1, 3p21.32 and 3p25.3-3p25.1. No significant correlation was found between the LOH and the grade of cancer differentiation. The difference between the LOH frequency in cervical carcinomas with early stage and those with advanced stage was not statistically significant. Only 51.4% of the 35 tumors were positive for HPV 16 E6 and 17.2% was HPV 18 E6 positive. HPV 16 was found to be positive in 64.7% of the Chinese, 41.7% in Malays and 50% in the Indians. Both the Malays and the Indians were observed to harbor the HPV 18 at a higher frequency (40 and 33.3% respectively) than the Chinese. In conclusion, besides HPV infection, other genetic abnormalities play a role in cervical carcinogenesis. LOH is a better method than chromosomal analysis in searching for possible tumor suppressor gene(s) that is responsible for cervical tumorigenesis. Mapping of the smallest region of LOH in these tumors and analysis of candidate genes present in the region of LOH will be continued.

Abstrak tesis yang dikemukakan kepada Senat Universiti Putra Malaysia sebagai memenuhi keperluan bagi mendapatkan Ijazah Doktor Falsafah

**PENYUSUNAN SEMULA KROMOSOM DAN KAJIAN KEHILANGAN
HETEROZIGOSITI DALAM BARAH SERVIKS DI KALANGAN PESAKIT
DI HOSPITAL KUALA LUMPUR**

Oleh

SITI NORLASIAH BINTI ISMAIL

Januari 2004

Pengerusi: Professor Abdul Manaf Ali, Ph.D

Institut: Biosains

Karsinoma serviks adalah malignan kedua yang kerap berlaku di kalangan wanita di seluruh dunia. Insiden yang paling tinggi kerap dilihat di negara yang sedang membangun. Populasi yang mempunyai insiden tinggi mungkin mudah mendapat karsinoma serviks akibat pendedahan terhadap papilomavirus manusia (HPV) dan gen yang diwarisi atau diperolehi. Infeksi HPV bukan selalu menyebabkan kanser serviks. Bagi kanser serviks, ciri genetik yang kerap dilihat adalah beberapa pengubahan genetik yang berulang-ulang, yang tak berkaitan dengan HPV. Jenis kerosakan genetik yang berulang-ulang adalah seperti amplifikasi gen, translokasi kromosom, delesi, kehilangan heterozigositi, mutasi titik, *transcriptional silencing* dan integrasi virus. Kajian ini dijalankan untuk mengetahui pengubahan gen di kalangan wanita yang mempunyai kanser serviks.

Kajian ini melibatkan 50 biopsi barah serviks dari wanita yang

menjalani rawatan barah serviks di Jabatan Obstetrik dan Ginekologi, Hospital Kuala Lumpur. Tujuan kajian ini adalah; i) membangunkan kultur sel primer daripada sel barah serviks bagi menentukan prevalen dan jenis aberasi kromosom, ii) pencirian kehilangan alel (*allelic loss*) pada kromosom 3p, 5p, 11 dan 17p (p53), seterusnya mengenalpastian lokasi gen penumpas barah (*tumor suppressor genes*) dan iii) untuk menentukan status HPV dalam kanser serviks. Kaedah kultur sel primer dan sitogenetik telah diperlakukan ke atas sampel biopsi barah serviks. Penjaluran-G telah digunakan untuk mengenalpasti kromosom. Untuk menentukan lokasi gen penumpas barah, kajian kehilangan heterozigositi telah dilakukan ke atas 37 sampel barah serviks. Dua puluh empat penanda polimorfik mikrosatelit pada kromosom 3p, 5p, 11 dan p53 telah dipilih. Pelet DNA normal dan barah dari setiap 37 sampel telah dianalisis menggunakan kaedah analisis mikrosatelit berasaskan PCR (*PCR-based microsatellite analysis*) untuk mengesan kehilangan alel.

Dua puluh lima biopsi barah serviks telah berjaya dikariotip dan *near-diploid* telah dilihat sebagai *modal number* dengan majoriti adalah hipoploid (35-45). Lebih kurang 50% daripada taburan metafasa adalah hipoploid, 12.1% hiperploid dan 36.7% adalah diploid. Keabnormalan pada jumlah kromosom dilihat lebih utama di kalangan pesakit, terutamanya monosomi kromosom 17, 22, X, 11, 18, 19, 13, dan 6. Penghibridan pendarfluor *in situ* (*fluorescence in situ hybridization*) menggunakan prob sentromerik bagi kromosom 11, 17 dan 18 telah

mengesahkan kehadiran monosomi kromosom tersebut tetapi pada kadar yang rendah, iaitu, 12.0%, 16.2% dn 26.4% masing-masing. Beberapa klon sel telah dijumpai, berkemungkinan 45,XX,-22; 45,XX,-11; 45,XX,-19 dan 45,XX,-18. Dua puluh enam daripada 36 individu yang informatif menunjukkan LOH pada satu lokus atau lebih. Insiden LOH yang tertinggi dikesan pada kromosom 3p dengan kekerapan pada kadar 48.6%, manakala LOH yang terendah adalah pada kekerapan 2.7% yang dilihat pada kromosom 17p di posisi 17p13.1. LOH telah dikesan pada empat kawasan yang utama, 11q23.3, 3p14.2-3p14.1, 3p21.32 dan 3p25.3-3p25.1. Tiada korelasi yang signifikan ditemui di antara LOH dan gred pembezaan kanser. Perbezaan di antara kekerapan LOH dalam karsinoma serviks di peringkat awal dan karsinoma di peringkat akhir adalah tidak signifikan. Hanya 51.4% daripada 35 barah serviks positif bagi HPV 16 E6 dan 17.2% positif untuk HPV 18 E6. HPV 16 didapati positif di dalam 64.7% kaum Cina, 41.7% Melayu dan 50% di kalangan kaum India. Peratusan positif HPV 18 didapati lebih tinggi di kalangan kaum Melayu dan India berbanding kaum Cina. Sebagai rumusan, selain infeksi HPV, keabnormalan genetik juga berperanan dalam proses karsinogenesis serviks. LOH adalah kaedah yang lebih baik daripada kaedah analisis kromosom untuk mencari gen penumpas barah yang bertanggungjawab bagi proses barah serviks. Kehadiran LOH dalam barah serviks mungkin boleh digunakan sebagai indikator terhadap prognosis penyakit tersebut. Pemetaan kawasan terkecil bagi LOH dalam barah ini serta analisis calon gen dalam kawasan LOH akan diteruskan.

بِسْمِ اللَّهِ الرَّحْمَنِ الرَّحِيمِ

ACKNOWLEDGEMENTS

First of all praise to the Almighty Allah for giving me the ability to learn and strength to complete this research. This study would never have materialized without the contribution and support of many people to whom I have the pleasure of expressing my appreciation and gratitude. I would like to express my sincere gratitude to Professor Dr. Abdul Manaf Ali for his guidance, understanding, patience and encouragement that lead to the completion of this thesis, and my supervisory committee members: Professor Datin Dr. Khatijah Mohd. Yusoff, for providing the facility for molecular work, monitoring the progress of research, helpful comments and editing of the thesis; Associate Professor Dr. Sabariah Abdul Rahman and Associate Professor Dr. Siti Aishah Mohd. Ali, for their invaluable comments and Dr. Noorjahan Banu Mohd. Ali-Theen for proofreading and editing this thesis.

I am greatly indebted to the full cooperation and support of many individuals in the Department of Obstetrics and Gynecology, Hospital Kuala Lumpur, for providing the samples, without them, this research would have not materialized. Thanks are extended to the ex-Director Dr. Alex Mathews and Dr. Wong Sum Keong and his medical and nursing team.

My sincere gratitude to the Director of Institute for Medical Research (IMR) and the staff of the Genetics Laboratory for allowing the use of their laboratory for the Interphase *in situ* hybridization technique. Special thanks to Dr. Puteri JamilatulNoor Megat Baharuddin for sharing her experience and knowledge in this technique and her continuous encouragement towards the completion of this postgraduate program, Ms. Roshidah Ishak for the valuable tips on PCR technique, Mrs. Wang Lily for her technical assistance in FISH technique, Mr. Roslaini Majid for the simplified Hematoxylin and Eosin (H & E) technique and Ms. Ten Sew Keoh for her input. I am grateful to Ms Subha Bhassu of the Department of Biochemistry and Microbiology, UPM for the microsatellite analysis technique. My sincere thanks to Dr. Norhayati Zainal Abidin of Malaya University for providing the CaSki cell line.

I would like to extend my sincere thanks to the staff of the Genetics Laboratory of the National Population and Family Development Board (NPFDB) for their technical assistance during the study period, including taking messages from the doctors for specimen collection. My gratitude to my organization, NPFDB, for allowing me to pursue the postgraduate program, and the Public Service Department, as on behalf for the Government of Malaysia, for sponsoring me throughout this program.

I greatly appreciated my colleagues from the Animal Cell Culture



Laboratory and Biochemistry Laboratory for their unduly support and encouragement during the hard times. Lastly, but not the least, to my husband and children for their patience, sacrifices, support and understanding throughout this postgraduate program, my beloved mother and my siblings for their unconditional love and encouragement throughout the trying period.



I certify that an Examination Committee met on 13th January 2004 to conduct the final examination of Siti Norlasiah binti Ismail on her Doctor of Philosophy thesis entitled "Chromosomal Rearrangement and Loss of Heterozygosity Studies in Cervical Cancer Amongst Patients in Hospital Kuala Lumpur" in accordance with Universiti Pertanian Malaysia (Higher Degree) Act 1980 and Universiti Pertanian Malaysia (Higher Degree) Regulations 1981. The Committee recommends that the candidate be awarded the relevant degree. Members of the Examination Committee are as follows:

Rozita Rosli, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Chairman)

Abdul Manaf Ali, Ph.D.

Professor
Institute Biosciences
Universiti Putra Malaysia
(Member)

Datin Khatijah Mohd Yusoff, Ph.D.

Professor
Faculty of Sciences and Environmental Studies
Universiti Putra Malaysia
(Member)

Sabariah Abdul Rahman, Ph.D.

Associate Professor
Faculty of Medicine and Health Sciences
Universiti Putra Malaysia
(Member)

Noorjahan Banu Mohd Ali-Theen, Ph.D.


Faculty of Food Science and Biotechnology
Universiti Putra Malaysia
(Member)

Sit Aishah Mohd Ali, M.D

Associate Professor
Faculty of Medicine
Universiti Kebangsaan Malaysia
(Member)

Mohd Nizam Hj. Musa, Ph.D.

Professor
International Medical University
(Independent Examiner)



GULAM RUSUL RAHMAT ALI, Ph.D.

Professor/Deputy Dean
School of Graduate Studies
Universiti Putra Malaysia

Date: 21 APR 2004

This thesis submitted to the Senate of Universiti Putra Malaysia has been accepted as fulfilment of the requirement for the degree of Doctor of Philosophy. The members of the Supervisory Committee are as follows:


Abdul Manaf Ali, Ph.D,
Professor,
Institute of Bioscience,
Universiti Putra Malaysia.
(Chairman)

Datin Khatijah Mohd Yusoff, Ph.D,
Professor,
Faculty of Environmental Sciences,
Universiti Putra Malaysia.
(Member)

Sabariah Abdul Rahman, M.D
Associate Professor,
Faculty of Medical Sciences,
Universiti Putra Malaysia.
(Member)

Noorjahan Banu Mohd Ali-Theen, Ph.D,
Faculty of Food Science and Biotechnology,
Universiti Putra Malaysia.
(Member)

Siti Aishah Mohd Ali, M.D,
Associate Professor,
Faculty of Medicine,
Universiti Kebangsaan Malaysia.
(Member)




AINI IDERIS, Ph.D,
Professor/Dean
School of Graduate Studies
Universiti Putra Malaysia.

Date: 17 MAY 2004

DECLARATION

I hereby declare that the thesis is based on my original work except for quotations and citations which have been duly acknowledged. I also declare that it has not been previously or concurrently submitted for any degree at UPM or other institutions.



SITI NORLASIAH BINTI ISMAIL

Date: 12 March 2004

TABLE OF CONTENTS

	Page
DEDICATION	ii
ABSTRACT	iii
ABSTRAK	vi
ACKNOWLEDGEMENTS	ix
APPROVAL	xii
DECLARATION	xiv
LIST OF TABLES	xviii
LIST OF FIGURES	xxi
LIST OF ABBREVIATIONS	xxiv
GLOSSARY	xxvi
CHAPTER	
I INTRODUCTION	1
II LITERATURE REVIEW	8
The multistep nature of cancer	8
Cervical cancer	11
Etiology of cervical cancer and its precursors	12
Classification and staging of cervical cancer	13
Human papillomavirus and cervical cancer	14
Screening and testing for HPV	20
Oncogenes in cervical cancer	21
Tumor suppressor genes	23
Genetic alterations in cervical cancer	27
Methods of localizing tumor suppressor genes	31
Cytogenetic studies of carcinoma of the cervix	32
Interphase cytogenetics of carcinoma of the cervix	40
Loss of heterozygosity and microsatellite instability in cervical cancer	42
Loss of heterozygosity (LOH) in cervical cancer	45
Microsatellite instability (MI/MSI) in cervical cancer	54
III MATERIALS AND METHODS	56
Sample population of women with cervical cancer	56
Clinical specimen of cervical tumor biopsies	56
Cytogenetic analysis of cervical cancer	57



	Primary culture	57
	Subculturing of monolayer cells by trypsinization	58
	Harvesting and slide preparation	59
	Banding and chromosome analysis	60
	Freezing of cells and storage	61
	Cell recovery	64
	Interphase cytogenetics by fluorescence <i>in situ</i> hybridization (FISH)	65
	Preparation of specimen and pretreatment of cells	65
	Prehybridization	66
	Post hybridization washing, counterstaining and visualization	67
	Loss of heterozygosity (LOH) or allelic loss analysis	70
	Preparation of genomic DNA	70
	Microsatellite polymorphisms	72
	Amplification of microsatellite loci	76
	Assessment of loss of heterozygosity (LOH)	78
	Detection of human papillomavirus DNA 16 and 18	78
	Culture of established cell lines	78
	Isolation of DNA from cultured cells	79
	Detection of Human Papillomavirus HPV DNA 16 and 18 sequences	79
IV	RESULTS AND DISCUSSION	83
	Sample population of women with cervical cancer	83
	Clinical specimen of cervical tumor biopsies	83
	Morphology of the original tumor	83
	Morphology of the cultured cells	88
	Cytogenetic analysis of cervical cancer	96
	Interphase Cytogenetics analysis of cervical imprints	112
	Loss of heterozygosity (LOH) or allelic loss analysis	123
	Loss of heterozygosity in the short-arm of chromosome 3	129
	Loss of heterozygosity in the short-arm of chromosome 5	140
	Loss of heterozygosity on chromosome 11	145
	Loss of heterozygosity at the region 17p13.1	152
	Detection of HPV DNAs 16 and 18	155
V	GENERAL DISCUSSION AND CONCLUDING REMARKS	162
	Further research	177

REFERENCES	179
APPENDICES	
Appendix A	206
Appendix B	211
Appendix C	224
BIODATA OF THE AUTHOR	229



LIST OF TABLES

Table	Page
2.1 1995 Modification of FIGO Staging of Carcinoma of the Cervix	15
2.2 Frequency of LOH in Cervical Cancer and SIL (CIN)	46
2.3 Incidence of allelic losses of markers on 3p	49
3.1 Definition of terms and recommendations related to abnormalities commonly seen in neoplasia	62
3.2 Microsatellite markers on 3p selected for the study	73
3.3 Microsatellite markers on 5p selected for the study	74
3.4 Microsatellite markers on 11p and 11q selected for the study	75
3.5 Parameters for polymerase chain reaction in the amplification of microsatellite polymorphisms on chromosome 3p, 5p, and 11	77
3.6 Sequences of oligonucleotide primer pairs for the detection of HPV 16 and 18	82
4.1 FIGO's classification of patients	84
4.2 Number of primary cell cultures successfully established according to clinical stages in the women diagnosed with cervical cancer	90
4.3 Number of successfully karyotyped cases according to clinical stages in 32 successfully established primary cultures	97
4.4 Cytogenetic findings in the twenty-five women diagnosed with cervical cancer	98
4.5 The incidence of chromosome abnormalities seen in the 25 patients diagnosed with cervical cancer	101



4.6	Distribution of successfully karyotyped cases according to age and type of chromosomal abnormalities	103
4.7	Clinical data and DNA ISH results for chromosome 11 in 23 women with cervical carcinoma	113
4.8	Clinical data and DNA ISH results for chromosome 17 in 23 women with cervical carcinoma	114
4.9	Clinical data and DNA ISH results for chromosome 18 in 13 women with cervical carcinoma	115
4.10	DNA ISH results for chromosome 11, 17 and 18 according to clinical stages	118
4.11	Loss of heterozygosity in 37 cervical carcinomas using 21 microsatellite polymorphic markers	124
4.12	Frequency of LOH with 21 microsatellite markers in 37 informative individuals with cervical carcinomas	126
4.13	Correlation of LOH with clinical characteristics in cervical carcinomas	126
4.14	Frequency of loss of heterozygosity at 9 loci on the short arm of chromosome 3 in 36 informative individuals with cervical carcinomas	131
4.15	Frequency of LOH according to clinical stages (FIGO) on the short arm of chromosome 3 in the informative individuals with cervical carcinomas	132
4.16	Frequency of loss of heterozygosity at 4 loci on the short arm of chromosome 5 in 37 patients with cervical carcinomas	141
4.17	Frequency of LOH according to clinical stages (FIGO) at the 4 loci on the short arm of chromosome 5 in 30 informative individuals with cervical carcinomas	142
4.18	Frequency of LOH at 7 loci on the chromosome 11 in 37 patients with cervical carcinomas	147

4.19	Frequency of LOH according to clinical stages (FIGO) at the 6 loci on the chromosome 11 in 37 informative individuals with cervical carcinomas	148
4.20	Frequency of LOH at locus for p53 in patients with cervical carcinomas	153
4.21	Frequency of LOH at 17p13.1 according to clinical stages (FIGO) in 11 informative individuals with cervical carcinomas	153
4.22	Outcomes of Human papillomavirus (HPV) 16 and 18 detection in 37 women with cervical carcinomas	156
4.23	The distribution of HPV 16 status according to clinical stage (FIGO) in 33 women with cervical carcinomas	157



LIST OF FIGURES

Figure	Page
2.1 Events in neoplastic transformation	10
2.2 Genome organization of HPV	17
3.1 Flow chart of the process of primary tissue culture and karyotyping	63
3.2 Overview of fluorescence <i>in situ</i> hybridization technique (FISH)	69
3.3 Protocol for DNA extraction from blood and tumor tissue	71
3.4 DNA extraction procedure for cultured cells	80
4.1 G4. Cervical imprint stained with H & E from the patient clinically diagnosed as having a stage IB1 cervical cancer	85
4.2 G19. Cervical imprint stained with H & E from the patient clinically diagnosed as having a stage IB cervical cancer	85
4.3 G9. Cervical imprint stained with H & E from the patient clinically diagnosed as having a stage IIA	86
4.4 G47. Cervical imprint stained with H & E from the patient clinically diagnosed as having a stage IIB	86
4.5 G5. Cervical imprint stained with H & E from the patient clinically diagnosed as having a stage IIIA	87
4.6 G15. Cervical imprint stained with H & E from the patient clinically diagnosed as having a stage IIIB	87
4.7 Cytological smear displaying cells suggestive of invasive squamous cell carcinoma	89



4.8	Cytological smear of a patient indicative of invasive squamous cell carcinoma	89
4.9	G4. Phase-contrast features of the monolayered cultured cells revealing a sheet of polygonal cells with a pavement-like arrangement (100X) cultured from a stage IB1 cervical tumor	91
4.10	G33. Phase-contrast features of the monolayered cultured cells revealing a sheet of polygonal cells with a pavement-like arrangement (100X) cultured from a stage IIB cervical tumor	91
4.11	Phase-contrast features of the established cervical cell line, CaSki cell, displaying a sheet of polygonal cells with a pavement-like arrangement (200X)	93
4.12	G11. Phase-contrast of the cultured cervical carcinoma exhibiting another type of cell morphology	93
4.13	G32. Phase-contrast microscopy of the cervical cell culture showing the presence of contamination of fibroblasts (100X)	94
4.14	G33. <i>In situ</i> hybridization of the interphase nuclei from cervical imprints	119
4.15	G37. <i>In situ</i> hybridization of the interphase nuclei from cervical imprints	119
4.16	G40. <i>In situ</i> hybridization of the interphase nucleus from cervical imprints	120
4.17	Patterns of LOH on 3p.	133
4.18	Ethidium bromide stained Metaphor™ gel of PCR products using primer flanking microsatellite polymorphism at D3S1228 locus on 3p.	134
4.19	Ideogram showing pattern of LOH on 5p using microsatellite polymorphisms in cancerous lesions of cervix.	143

4.20	Ideogram showing pattern of LOH on chromosome 11 in cancerous lesions of cervix.	146
4.21	Ethidium bromide stained agarose gel (1.5%) of PCR products of cervical cancer patients tested for HPV type 16 E6	157



LIST OF ABBREVIATIONS

CIN	Cervical intraepithelial neoplasia
CGH	Comparative genomic hybridization
dATP	Deoxyadenosine triphosphate
dCTP	Deoxycytidine triphosphate
dGTP	Deoxyguanosine triphosphate
DMSO	Dimethylsulfoxide
dTTP	Deoxythymidine triphosphate
FISH	Fluorescence <i>in situ</i> hybridization
G-band	Giemsa band
HCl	Hydrogen chloride
H & E	Hematoxylin and Eosin
HPV	Human papillomavirus
ISCN	International System for Human Cytogenetic Nomenclature
ISH	<i>In situ</i> hybridization
KCl	Potassium chloride
LOH	Loss of heterozygosity
mar	Marker chromosome
MOH	Ministry of Health
N/C	Nucleus and cytoplasmic ratio
NPFDB	National Population and Family Development Board
PBS	Phosphate buffered-saline
PCR	Polymerase chain reaction

